



**Universitat de Lleida**

Document downloaded from:

<http://hdl.handle.net/10459.1/60215>

The final publication is available at:

<https://doi.org/10.1016/j.foreco.2017.07.039>

Copyright

cc-by-nc-nd, (c) Elsevier, 2017



Està subjecte a una llicència de [Reconeixement-NoComercial-SenseObraDerivada 4.0 de Creative Commons](https://creativecommons.org/licenses/by-nc-nd/4.0/)

1    **Is silviculture able to enhance wild forest mushroom resources? Current knowledge and**  
2    **future perspectives**

3

4    Antonio Tomao<sup>a\*</sup>, José Antonio Bonet<sup>b,c</sup>, Juan Martínez de Aragón<sup>c</sup>, Sergio de-Miguel<sup>b</sup>

5    <sup>a</sup>Department for Innovation in Biological, Agro-food and Forest systems (DIBAF), University of Tuscia, Via S. Camillo  
6    de Lellis, s.n.c. -I- 01100, Viterbo, Italy. Tel. +39 0761357407 Fax: +39 0761357389

7    <sup>b</sup>Department of Crop and Forest Science, University of Lleida-Agrotecnio Center (UdL-Agrotecnio), Avda. Alcalde  
8    Rovira Roure, 191, E-25198 Lleida, Spain

9    <sup>c</sup>Forest Sciences Centre of Catalonia (CTFC-CEMFOR), Ctra. de St. Llorenç de Morunys, km 2, E-25280 Solsona,  
10    Spain

11    \*Corresponding author: antonio.tomao@unitus.it

12

13    **Abstract**

14    Fungal fruit-bodies are an important economic resource both for recreational pickers and commercial  
15    markets. The broad interest in forest fungi and mushrooms has motivated attempts to develop  
16    appropriate silvicultural methods able to preserve and improve mushroom yields. Defining best  
17    practices for the sustainability and profitability of forest fungal resources is the main aim of  
18    ‘mycosilviculture’. However, the difficulty of monitoring and studying such a cryptic kingdom (i.e.,  
19    fungi) under real forest conditions has led to rather scattered scientific knowledge of the effects of  
20    forest management regimes and silvicultural operations on wild mushroom resources. Here, we  
21    review the current scientific literature regarding the influence of (i) stand characteristics, i.e. stand  
22    age, stand density, canopy cover and tree species composition, (ii) silvicultural practices and (iii)  
23    other management-related disturbances affecting the yield of wild epigeous mushrooms, with the aim  
24    of systematizing existing scientific knowledge and identifying gaps in knowledge in order to suggest

25 future lines of research. Most of the research in the field of mycosilviculture to date has focused on  
26 ectomycorrhizal mushrooms, which include the majority of wild commercial mushrooms. The main  
27 findings from the literature indicate that forest management practices, by modifying stand  
28 characteristics and microclimatic conditions, can influence wild mushroom occurrence and  
29 productivity, both positively and negatively, depending on the specific fungal ecological needs,  
30 reproductive strategies, forest type and management regime. Further research efforts relating to all  
31 aspects of the interaction between forest management, fungal ecology and mushroom production are  
32 needed: in particular, additional research efforts should be devoted to understanding mushroom yield  
33 dynamics in uneven-aged and mixed forests and evaluating the effects of regeneration methods on  
34 fungal communities.

## 35 **Keywords**

36 Mycosilviculture; yield; mushrooms; forest management; fungi.

## 37 **1. Introduction**

38 Fungi are widely recognized as fundamental components of biodiversity and ecosystem functioning.  
39 Furthermore, fungal fruit-bodies, which constitute the main resource of an important socioeconomic  
40 activity based on mushroom picking for both recreational purposes and commercial markets, provide  
41 important provisioning and cultural ecosystem services to society. The commercial value of  
42 mushrooms can exceed 40 million euros (€) in countries such as Italy (ISTAT, 2008) or 32 million €  
43 in regions such as Catalonia (Bonet et al., 2014). The value derived from mushrooms is even higher  
44 if we also consider the recreational value of mushroom picking as a leisure activity (Martínez de  
45 Aragón et al., 2011; Latorre, 2016). Indeed, the economic profit from wild mushrooms can be higher  
46 than timber revenues (Palahí et al., 2009; Pettenella and Secco, 2006), especially in areas where  
47 timber production is not very profitable. Furthermore, the economic profit from wild mushrooms can  
48 represent a considerable percentage (up to 25%) of the soil expectation value, even in countries such

49 as Finland where timber production is an important economic activity (Tahvanainen et al., 2016). De  
50 Roman and Boa (2006) calculated that a family of four people living in a rural area of Northern Spain  
51 can derive profits of 5,600–8,400 € from picking saffron milk caps (*Lactarius deliciosus s.l.*) over the  
52 course of a season. In Eastern Finland, Cai et al. (2011) reported that the seasonal earnings of  
53 professional pickers were 1,224 euro (more than 5% of the average net annual household income in  
54 Finland) and more recently Sisak et al. (2016) reported that three quarters of the rural households in  
55 the Czech Republic collect Non Wood Forest Products (NWFPs) (mainly mushrooms and berries).  
56 However, quantifying the exact contribution of mushrooms to household incomes in economic terms  
57 may still be underestimated due to the difficulty in accurately estimating the amount of wild  
58 mushrooms that are sold (because the economy is largely informal) (Maso, 2008).

59 The twofold importance of wild forest mushrooms in ecological and socioeconomic terms underlines  
60 the need for promoting and preserving this resource (Dettori et al., 2009). For these reasons, forest  
61 management regimes oriented toward the provision of wild mushrooms need to be based on reliable  
62 quantifications of the impact of anthropogenic disturbances on mushroom yields, which also implies  
63 an in depth scientific knowledge of the key drivers that affect fungal communities.

64 Sporocarp production has been the focus of many studies (Vogt et al., 1992) and numerous variables  
65 influencing mushroom production have been identified. In addition to specific plant–fungal  
66 interactions, other key factors affecting sporocarp emergence include climatic conditions and soil and  
67 topographical characteristics (Figure 1). Water availability is fundamental for fruit-body formation  
68 (Wiklund et al., 1995; Ogaya and Peñuelas, 2005; Büntgen et al., 2012). In particular, mean monthly  
69 precipitation and accumulated mean monthly evapotranspiration are strongly related to annual fruit-  
70 body production (Martínez de Aragón et al., 2007; Taye et al., 2016). Climatic conditions may also  
71 affect mushroom production in the long term (Gange et al., 2007; Kauserud et al., 2008; Ágreda et  
72 al., 2016) as a consequence of climate change. However, climatic or site conditions alone do not  
73 completely explain mushroom occurrence (Barroetaveña et al., 2008; Egli et al., 2011; Krebs et al.,

2008) because the interactions between individual meteorological variables and other ecosystem variables are quite complex (Egli et al., 2011).

Stand structure characteristics (e.g., tree species composition, stand age, and stand density) influence mushroom yields (Figure 1). Identifying stand characteristics that enhance the productivity of edible and marketable wild mushrooms is also the basis for the development of decision-support tools based on predictive models, which are being used to select optimal forest management alternatives for the provision of multiple ecosystem services, e.g., the joint production of mushrooms and timber (Palahí et al., 2009; Tahvanainen et al., 2016).

(FIGURE 1 HERE)

The growing interest in mushroom production has led to the development of a promising new discipline over the past few decades: “mycosilviculture”. According to Martínez-Peña et al. (2011), mycosilviculture attempts to integrate timber and mushroom production into the management of forest ecosystems. However, a broader, more generalized definition of mycosilviculture based on existing formal definitions of silviculture (e.g., Ciancio, 1981; Ciancio and Nocentini, 1997) is: the experimental science studying the interactions between the natural dynamics of forest ecosystems and management techniques, with the aim of defining the best practices for the sustainability and profitability of fungal resources.

Although the best practices necessary for truffle (*Tuber* spp.) cultivation are well established (e.g., Bonet et al., 2009; Giovannetti et al., 2010; Reyna and Garcia-Barreda, 2014), this is not the case for epigeous macromycetes, some of which are also commercially important. Most of the economically valuable species are mycorrhizal fungi (e.g. *Boletus* spp., *Lactarius* spp., *Cantharellus* spp. and *Tricholoma* spp.). However, some saprotrophic (e.g. *Morchella* spp., *Agaricus* spp. and *Macrolepiota*

spp.) and parasitic (e.g. *Armillaria* spp.) species may also be important for local economies (Table 1). Although an increasing number of studies have focused on evaluating the effects of silvicultural practices on fungal yield in recent years (e.g., Bonet et al., 2012; Lin et al., 2015), and some authors have proposed guidelines for fungal-oriented silviculture (Palahí et al., 2009; Pierangelo and Rolland, 2013; Bettini et al., 2016), the large number of potential variables influencing mushroom yields, and their interdependence, make the development of clear forest management recommendations rather challenging. Moreover, the existing studies are mostly local or regional and, hence, the scientific knowledge on this topic is rather scattered. Therefore, recommendations that may be useful for a specific area may not be suitable for sites with a different climate or forest stand structure, or if the highly valued edible mushroom species differ between regions according to differences in stakeholders' preferences. In this sense, a systematic review of current knowledge about forest management practices that are able to increase mushroom yield may help to shed light on this complex issue and contribute to clarifying the state of the art as a basis for further research.

111

(TABLE 1 HERE)

113

Based on these considerations, the aims of this paper are: (i) to review the current scientific literature regarding the effects of forest management on the yield of wild epigeous mushrooms, with a particular focus on those that are mainly used worldwide as a food provisioning ecosystem service and that are also linked to the provision of cultural ecosystem services (recreation and tourism) (Table 1); and (ii) to identify gaps in our knowledge and suggest new research lines.

Following this introduction, Section 2 reviews the stand variables that are known to affect mushroom production, i.e., forest age, stand density, canopy cover and tree species composition. Section 3 deals with the forest management practices that may influence wild epigeous mushroom occurrence by

122 modifying stand characteristics and microclimatic conditions i.e., the effect of thinning, regeneration  
123 methods and other management practices and anthropogenic disturbances (i.e., logging, mushroom  
124 picking, trampling, raking, shrub control and prescribed fire). Lastly, Section 4 summarizes the main  
125 findings from the literature review, discusses gaps in our knowledge and recommends areas for future  
126 research.

## 127 **2. Stand structure variables influencing mushroom production**

### 128 **2.1 Forest stand age**

129 Studies have shown that forest age is an important factor in mushroom fruiting abundance. However,  
130 the ecology of fungal species can vary considerably even within the same genus and, therefore,  
131 providing a general overview of the results is difficult. Indeed, wild mushrooms can be classified as  
132 typical of either early- or late-stage forest succession (Mason et al., 1987) according to their  
133 physiology and nutritional requirements (Dighton and Mason, 1985), although some species can be  
134 present in both early and late successional stages: e.g., *Suillus granulatus* in pine forests (Savoie and  
135 Largetau, 2011). Furthermore, the concept of an 'old' or 'young' stand may depend, e.g., on the fertility  
136 of the site or the longevity of the tree species (Martínez-Peña et al., 2012a). For these reasons,  
137 mushroom fruiting can vary depending on forest type and fungal species.

138 Many studies agree that sporocarps of mycorrhizal fungi are more abundant in younger stands. This  
139 evidence has been related to the physiological status (higher growth rate) of trees in early stage stands  
140 (Senn-Irlet and Bieri, 1999; Bonet et al., 2008; Egli et al., 2010; Tahvanainen et al., 2016). For  
141 example, sporocarp production in young and open *Picea excelsa* stands was almost double that of  
142 older stands (Senn-Irlet and Bieri, 1999). In a study conducted in mixed-coniferous forests in Russia,  
143 Chibisov and Demidova (1998) reported that mushroom production was greatest in young stands (15–  
144 35-years-old). In Finland, a study on mushroom yields in pine forests of different ages reported that  
145 the 20–30-year-old stands produced greater quantities of sporocarps than the other stands (Hintikka,

146 1988). Similar results have been found by Kalamees and Silver (1988) in *Pinus sylvestris*-dominated  
147 stands, where both total and edible mushroom yields were greater in 25-year-old stands than in other  
148 stands. Smith et al. (2002) found that the average epigeous sporocarp yield in *Pseudotsuga menziesii*  
149 forests in Oregon (USA) was about six times lower in very old (more than 400 years old) stands  
150 compared with young and rotation-age stands.

151 Similar findings have been reported regarding the productivity of important edible, marketable  
152 species such as *Lactarius deliciosus* s.l.. Martínez de Aragón et al. (2007) reported that *Lactarius*  
153 *sanguifluus* sporocarps were more abundant in young forest stands (<50 years old) and that although  
154 stand age was not correlated with total sporocarp yield, it was negatively associated with the amount  
155 of saffron milk caps. This group of species is mainly found in conifer stands that are less than 40  
156 years old (Fernández-Toirán et al., 2006; Kranabetter et al., 2005; Kucuker and Baskent, 2015).  
157 Furthermore, the results of a 15-year study monitoring mushroom production in *Pinus pinaster* stands  
158 in Central Spain revealed that saffron milk caps were found in all age classes, but that significantly  
159 higher yields were obtained in the youngest stands (11–20 years old) (Ágreda et al., 2014). In a similar  
160 study, Martínez-Peña et al. (2012a) reported that the *L. deliciosus* sporocarp yield peaks twice in  
161 stands dominated by Scots pine: the first peak in yield occurs in stands that are between 16 and 30  
162 years old and then again when the stand is more than 71 years old. The authors suggested that these  
163 findings reflect the preference of this species for open-canopy conditions, which can also be found  
164 within mature stands in intensively managed forest ecosystems.

165 The influence of stand age on the production of *Boletus edulis* s.l. has also been studied. Keizer and  
166 Arnolds (1994) reported that this species group was found more frequently in 20- to 40-year-old  
167 *Quercus* and *Fagus* forests than in younger (i.e., 10–20-year-old) and older (50–140-year-old) forests.  
168 Higher yields of *B. edulis* have also been reported in mature pine stands (Hintikka, 1988; Keizer and  
169 Arnolds, 1994). For example, Martínez-Peña et al. (2012a) found that *B. edulis* sporocarp production  
170 was greater in 51–70-year-old *Pinus sylvestris* stands than in younger stands. In Norway spruce



171 (*Picea excelsa*) stands, *B. edulis* sporocarp production reaches its highest levels 25–30 years after  
172 planting, just before the first thinning usually takes place (Tahvanainen et al., 2016).

173 Ortega-Martínez et al. (2011) evaluated the effect of forest age on the weekly growth of sporocarps,  
174 and reported that both *B. edulis* and *L. deliciosus* fruit-bodies grow faster when they are associated  
175 with young Scots pine trees (<30-years-old). Sporocarps were larger and nearly 50% heavier  
176 compared with mushrooms of the same species growing in older pine stands, which may be because  
177 young *Pinus sylvestris* trees require more nutrients to support their rapid growth rates during the early  
178 stand development stage, and the ectomycorrhizal fungi may benefit from the higher carbohydrate  
179 allocation to the root system. By contrast, American Matsutake (*Tricholoma magnivelare*) is  
180 generally found in mature stands that are more than 70 years old (Kranabetter et al., 2002; 2005).

## 181 **2.2 Forest stand density and canopy cover**

182 The composition of the mycorrhizal fungal community is strongly influenced by forest stand structure  
183 and density owing to their close symbiotic relationship with trees (Santos Silva et al., 2011). Stand  
184 basal area is one of the most used variables to describe the density of forest stands and is a strong  
185 predictor of fungal yield (e.g., Bonet et al., 2008; 2010; de-Miguel et al., 2014; Tahvanainen et al.,  
186 2016).

187 In *Pinus sylvestris* stands in the Central Pyrenees, Bonet et al. (2008) found that maximum mushroom  
188 yield was achieved under stand basal areas of approximately 20 m<sup>2</sup> ha<sup>-1</sup>. Similar findings have been  
189 reported from a study based on 21 plots established in *Pinus halepensis*, *Pinus nigra* and *Pinus*  
190 *syvestris* stands in Northeastern Spain (Bonet et al., 2010): maximum productivity of total, edible  
191 and marketable mushrooms was recorded in low–medium density stands with stand basal areas  
192 ranging from 15 to 20 m<sup>2</sup> ha<sup>-1</sup>. More recently, de-Miguel et al. (2014) reported that the optimal stand  
193 basal area that maximizes edible and marketable mushroom yield may differ among Mediterranean  
194 pine forest ecosystems (i.e., pure *P. sylvestris*, *P. nigra*, *P. halepensis* and *P. pinaster* stands, and

195 mixed *P. sylvestris*–*P. nigra* and *P. nigra*–*P. halepensis* stands), ranging between 10–15 m<sup>2</sup> ha<sup>-1</sup> in  
196 pure *P. halepensis* stands (growing under rather arid conditions) and 35–40 m<sup>2</sup> ha<sup>-1</sup> in pure *P. pinaster*  
197 stands growing under good conditions.

198 Martínez-Peña et al. (2012b) determined that stand basal area affects *Boletus edulis* yield, with  
199 productivity predicted to peak in the *P. sylvestris* forests of Central Spain when the stand basal area  
200 is approximately 40 m<sup>2</sup> ha<sup>-1</sup>. Tahvanainen et al. (2016) have also identified the optimal stand basal  
201 area that maximizes the production of *B. edulis* (approximately 25 m<sup>2</sup> ha<sup>-1</sup>) and *Lactarius* spp.  
202 (approximately 30 m<sup>2</sup> ha<sup>-1</sup>) in Norway spruce stands in Finland.

203 Stand density can also be described by the number of trees per hectare. However, this alternative  
204 quantification of density seems to be less relevant than stand basal area for understanding mushroom  
205 occurrence and yield. For example, Kucuker and Baskent (2015) found that this variable was not  
206 useful for predicting the occurrence of two *Lactarius* species (*L. deliciosus* and *L. salmonicolor*) in  
207 the Kızılcaasu Planning Unit, Turkey.

208 Canopy cover is also linked to forest stand density, and its effect on fungal yield has been studied in  
209 various forest ecosystems. Stands without canopy closure seem to be more suitable for mushroom  
210 production, especially in the case of important edible and marketable species such as *Lactarius*  
211 *deliciosus* s.l. and *Boletus edulis* s.l. For example, Ágreda et al. (2014) reported that *L. deliciosus*  
212 benefits from open stands because mushroom yields were higher in young forests without canopy  
213 closure than in forests with canopy closure. Martínez-Peña et al. (2012a) also reported higher  
214 *Lactarius* yields in open stands – both young and old – than in closed stands. These findings were  
215 supported by a Turkish study (Kucuker and Baskent, 2015): if canopy closure increases, the  
216 probability that *L. deliciosus* and *L. salmonicolor* sporocarps will be absent also increases. These  
217 authors hypothesized that canopy closure has an effect on soil temperature and moisture. Even *Boletus*  
218 *edulis* may benefit from stands without total canopy closure (Salerni and Perini, 2004). Likewise, in

219 *Cistus ladanifer* shrublands, a canopy cover of, for example, 80% supports sporocarp emergence  
220 better than total closure (Hernández-Rodríguez et al., 2015a).

### 221 **2.3 Tree species composition**

222 There is a strong relationship between macrofungal communities and vascular plant composition  
223 (e.g., Barluzzi et al., 1992; Packham et al., 2002; Landi et al., 2015), and the effect of tree species  
224 composition on the fungal community can be stronger than the influence of soil properties (Urbanová  
225 et al., 2015). This is partly due to the preference of some ectomycorrhizal species for specific host  
226 trees, as well as to differences in the litter properties and tree debris (i.e., needles, leaves, and cones)  
227 in the case of saprotrophic fungi (Zhou and Hyde, 2002).

228 Fungal species can be either generalists or specialists, depending on their ability to establish  
229 mycorrhizal associations with different tree species under a wide range of growing conditions or only  
230 with a given tree species under a particular range of environmental conditions, respectively. In the  
231 Holarctic realm, most ectomycorrhizal fungi are usually associated with several host trees (Kennedy  
232 et al., 2003). However, some important exceptions have been identified. *Suillus* and *Rhizopogon*  
233 genera are *Pinaceae*-specific fungi (Molina and Trappe, 1982). Furthermore, there are many  
234 symbionts that are exclusively associated with *Alnus* tree species (Molina, 1979; Tedersoo et al.,  
235 2009). Host specificity may be found not only at the genus level but also at the species level. For  
236 example, *Lactarius deterrimus*, which show a clear preference for Norway spruce under temperate  
237 conditions (Buée et al., 2011). Among important marketable species, *L. deliciosus* s.l. forms  
238 mycorrhizae with several *Pinaceae* species, whereas *Boletus edulis* and *Cantharellus* spp. are more  
239 generalist species that may establish symbioses with several coniferous and broadleaf tree species at  
240 different latitudes and altitudes. Some of the tree species associated to these prized marketable  
241 mushrooms are commonly used in reforestation (e.g. *Pinus*, *Picea*, *Pseudotsuga*, *Abies*) in the  
242 northern hemisphere and can produce relevant revenues from timber. This facilitates co-management  
243 of timber and mushrooms.

244 While it is accepted that fungal diversity increases with increasing tree species richness (Landi et al.,  
245 2014), the effect of tree composition on mushroom production is less clear. In this regard, Weigand  
246 (1997) reported that the productivity of *Tricholoma magnivelare* can be enhanced by between 100%  
247 and 400% by changing the tree composition and increasing the presence of American Matsutake host  
248 trees. In a long-term experiment in mixed forests, Egli et al. (2010) showed that the productivity of  
249 host-specific epigeous mushrooms is strongly related to the relative abundance of the host tree, which  
250 can be influenced through silviculture.

### 251 **3. Effect of forest management practices**

#### 252 **3.1 Thinning**

253 Thinning is the removal of a number of trees within a stand with the aim of enhancing the growth of  
254 the remaining trees by reducing the competition between them. As a man-induced disturbance of  
255 forest ecosystems, it may affect biodiversity and other related ecosystem functions (Bengtsson et al.,  
256 2000), including fungal communities (Lin et al., 2015), especially if combined with other  
257 environmental drivers (Tedersoo et al., 2011) such as climate factors (Kausarud et al., 2008). Since  
258 thinning modifies the humidity and light penetration under the canopy, thinned stands may experience  
259 more accelerated wetting and drying of the forest soil (Pilz and Molina, 2002). This can lead to a  
260 change in the presence and abundance of fungal species based on their specific temperature and water  
261 requirements. Furthermore, by removing the associated host trees, thinning may also eliminate some  
262 mycorrhizal fungi that were colonizing tree roots. At the same time, several mycorrhizal fungi  
263 associated to released trees may thrive (Bonet et al., 2012). The increased growth of the remaining  
264 trees may result in an increased allocation of carbohydrates to the mycorrhizal fungi, which, in turn,  
265 may partly aid them in the production of carpophores, i.e., healthy stands have a tendency to produce  
266 more mushrooms of ectomycorrhizal fungal species than weak and slow-growing trees (Savoie and  
267 Largetau, 2011). Therefore, the net effect of thinning on mycorrhizal mushroom yields, at least in the

268 short-term, is the result of the trade-off between immediate reduction in colonies and increased  
269 production of remaining mycorrhizal fungi.

270 In addition, there may be an increase in fruitbody production of non-mycorrhizal species after  
271 thinning, including both saprotrophic-terricol and saprotrophic-lignicolous species, as shown by Egli  
272 et al. (2010) in a long-term experiment in Switzerland. However, Kim et al. (2010) reported no  
273 difference in the occurrence of *Armillaria* spp., a very common and appreciated edible group of  
274 parasitic species, in thinned stands compared to unthinned stands.

275 The influence of thinning on fruitbody abundance can even be seen the first year after thinning (Bonet  
276 et al., 2012), although various studies have reported that thinning can have very different effects (i.e.  
277 either positive or negative) on mushroom yields. Normally, mushroom production seems to decrease  
278 after the disturbance (Pilz et al., 2006; Egli et al., 2010). However, Bonet et al. (2012) showed that  
279 *Lactarius deliciosus* s.l. yields were higher in thinned stands than in unthinned stands from the first  
280 productive season after the silvicultural cut, probably due to the low impact of the forest harvest  
281 operations in the forest area. Kropp and Albee (1996) also reported a similar trend for the same  
282 mushroom species in *Pinus contorta* stands and Tahvanainen et al. (2016) suggested that thinning  
283 would promote *B. edulis* yields in spruce plantations in Finland. A significantly positive correlation  
284 between fruitbody production (number of sporocarps) and tree-ring width was also shown in a thinned  
285 mixed forest (Egli et al., 2010) and in a dry *Pinus pinaster* ecosystem (Primicia et al., 2016).

286 However, some authors have reported that thinning has a negative impact on mushroom resources,  
287 with a decline in sporocarp production regardless of tree removal intensity, even if this effect varied  
288 by season (Luoma et al., 2004). Pilz et al. (2006) also found that the number and biomass of a  
289 chanterelle species (*Cantharellus formosus*) significantly decreased in the first year after thinning,  
290 even if no differences were observed in the mid-term (six years after treatment).

291 These contradictory results may be explained by the fact that thinning does not have the same effect  
 292 on all species. Effects are adverse or positive depending on the specific ecological needs of different  
 293 fungal species, their reproductive strategies and forest type (Pilz and Molina, 2002; Buée et al., 2005).  
 294 Kropp and Albee (1996) demonstrated that species of the genus *Suillus* (*Suillus brevipes*, *Suillus*  
 295 *tomentosus*) appear to be favored by changes resulting from silvicultural practices. *L. deliciosus* also  
 296 benefits from open stands, so thinning can positively influence its productivity (Bonet et al., 2012).  
 297 However, certain groups of ectomycorrhizal fungi (in particular the Hygrophoraceae) are negatively  
 298 affected by even low-intensity thinning (Kropp and Albee, 1996). The productivity of *Cantharellus*  
 299 species, which requires wet conditions to produce fruitbodies, seems to be reduced in fast-drying soils  
 300 after thinning (Pilz et al., 2006). After a silvicultural intervention in a 12-year-old Scots pine  
 301 plantation, Shaw et al. (2003) reported that only three ectomycorrhizal species (*Suillus bovinus*,  
 302 *Gomphidius roseus* and the not-edible *Cortinarius semisanguineus*) out of 19 considered in the study  
 303 showed an increase in sporocarp production.

304 Thinning may also influence the availability of host trees for target species (Landi et al., 2015) by  
 305 selecting particular tree species: preserving host trees that are suitable for target species may increase  
 306 the production of mushrooms, as demonstrated by Weigand (1997) for *Tricholoma magnivelare*.

307 Thinning intensity often varies according to different silvicultural objectives and can influence  
 308 mushroom production. Various levels of thinning intensity can be identified based on the quantity of  
 309 trees removed. Thus, in light thinning, the reduction in the stand basal area is usually less than 15–  
 310 20%, while in high-intensity thinning, the stand basal area may be reduced by more than 35–40%  
 311 (Mäkinen and Isomäki, 2004). Medium-intensity thinning has been reported to be preferable to  
 312 improve mushroom yields. Salerni and Perini (2004) observed that a greater amount of *B. edulis*  
 313 sporocarps were obtained from forest stands undergoing moderate thinning (removal of  
 314 approximately 20% of the stand basal area). Egli and Ayer (1997) demonstrated that a 35% reduction  
 315 of stems in a mixed forest in Switzerland increased edible mushroom production up to six-fold.

316 Conversely, heavy thinning, characterized by basal area removal ranging from 40% up to  
317 approximately 75%, seemed to inhibit *B. edulis* (Salerni and Perini, 2004) and *Cantharellus formosus*  
318 production (Pilz et al., 2006). This negative effect of heavy thinning may be related to the twofold  
319 effect of an accelerated drying of forest soils and the mortality of many colonies of mycorrhizal fungi  
320 so that the net effect on mushroom yields is consistently negative. Bonet et al. (2012) also studied the  
321 effect of thinning intensity on mushroom yields in Mediterranean pine forests, and found that light  
322 thinning was more beneficial than heavy thinning in terms of the yields obtained for the target group  
323 of fungal species (*L. deliciosus s.l.*). Maximum mushroom yields were found when stand basal area  
324 removal was approximately 10 m<sup>2</sup> ha<sup>-1</sup>. Conversely, heavy thinning can have negative effects on  
325 mushroom yields (Bonet et al., 2012).

### 326 **3.2 Regeneration methods**

327 Regeneration methods are harvesting techniques that aim to successfully establish a new cohort of  
328 trees that contributes to forest regeneration. Regeneration methods can be categorized into three main  
329 groups: clear-cutting, shelterwood methods and selective cuts. On the one hand, clear-cuts and  
330 shelterwood methods are typical of even-aged forestry, where successive thinnings may be combined  
331 during the rotation period with the possibility of performing a final cut in which the tree cover of  
332 adult trees is removed at the end of the rotation. On the other hand, selective cutting is the typical  
333 regeneration method in uneven-aged forests, also called continuous cover forestry, where tree cover  
334 is continuously maintained (Duryea and Dougherty, 2012; Pukkala and von Gadow, 2012).

335 Clear-cutting is a forestry practice in which all the trees of a given area are cut down. It has been  
336 reported to negatively influence the fruiting of most edible ectomycorrhizal fungi for approximately  
337 a decade until a new cohort of host trees establishes and their growth rate is sufficient to provide the  
338 ectomycorrhizal fungi with enough carbohydrates to support mushroom sporocarps (Kardell and  
339 Eriksson, 1987; Ohenoja, 1988; Durall et al., 2006). This effect has been reported even if the soil of  
340 a clear-cut area contains a quantity of potential inoculum similar to the adjacent forest area (Harvey

et al., 1980; Le Tacon, 1997). Furthermore, Parladé et al. (2017) reported that clear-cutting also has a negative effect on the survival of mycelium in the soil. By contrast, Hintikka (1988) found that mushroom yields recovered in the first years after the regeneration cut, when young and vital specimens are growing. Clear-cutting is also widely used for coppice management. Lee et al. (2016) found that the presence of *Armillaria* spp. increased in the first seven years after clear-cutting mixed oak stands. This result depends on the ecology of *Armillaria mellea* s.l., which includes both saprotrophic and pathogenic species that mainly infect trees with compromised physiology.

Shelterwood methods are another silvicultural option that can be used to enhance stand regeneration. They can be regarded as a progression of high-intensity thinning by preserving some standing, living, adult trees that will produce and spread seeds for the establishment of a new cohort of trees. Mushroom production by mycorrhizal fungi is higher in stands that were regenerated using shelterwood methods than in clear-cut stands because the remnant trees act as a fungal reservoir from which mycorrhizal fungi colonize the roots of the new cohort of trees (Peter et al., 2013). Similar results have been reported by Ágreda et al. (2014), who monitored fungal production in a chronosequence in *P. sylvestris* forests in Central Spain. They found that edible fungi were less abundant in the 10 years after shelterwood regenerative cuts. However, the presence of seed trees helped to support mushroom production. For this reason, the authors suggest the maintenance of seed trees as a relevant silvicultural tool to ensure the production of sporocarps and tree regeneration.

Selective cuts refer to small-scale forest management measures that aim to remove either single trees or small groups/patches of trees in order to facilitate regeneration in the resulting forest gap (Pukkala and Von Gadow, 2012). These operations have been found to influence fungal communities: the production of ectomycorrhizal mushrooms can be lower in the gap created by the selective cut compared to the surrounding uncut stands (Grebenc et al., 2009), particularly if all the trees in the gap are removed (De Groot et al., 2016).



365 However, the effect of regeneration cuts may differ among mushroom species, based on their ability  
366 to colonize a particular site after natural or man-induced disturbances. Fungi with resistant propagules  
367 or a high capacity for spore dispersal may rapidly colonize a site. Conversely, other species need an  
368 undamaged mycorrhizal network to connect them to the new host tree (Peter et al., 2013).  
369 Furthermore, differences in colonization ability are related to the different strategies used by fungi in  
370 the competition for water and nutrient uptake versus other fungal species.

### 371 **3.3 Use of mycorrhized trees**

372 Techniques for inoculating tree seedlings with ectomycorrhizal fungi have been widely studied;  
373 however, these studies have mainly focused on ectomycorrhizal fungi that promote tree growth and  
374 do not produce edible mushrooms or are not valued as food (Le Tacon et al., 1992; Savoie and  
375 Largetoie, 2011).

376 With regard to edible mushrooms, mycorrhized trees are successfully used to establish forest  
377 plantations for producing hypogeous edible fungi (truffles); however, these practices are more related  
378 to arboriculture than silviculture. Mycorrhized trees are not used to produce edible epigeous  
379 mushrooms in natural or planted forest ecosystems owing to a lack of evidence that this practice  
380 enhances the production of valuable ectomycorrhizal mushrooms. However, some promising results  
381 were observed by Martínez de Aragón et al. (2012) who tested the viability of introducing *Quercus*  
382 *ilex* seedlings inoculated with *Tuber melanosporum* (a hypogeous edible fungal species) into forest  
383 areas affected by large forest fires. The presence of the associated target mycorrhizae was observed  
384 in the seedlings ten years after planting. Unfortunately, no similar studies using wild edible epigeous  
385 mushrooms seem to have been performed.

386 Among the huge diversity of fungal species, only a small number of species have been domesticated.  
387 Currently, commercial nurseries offer a limited variety of mycorrhized plants inoculated with  
388 marketable fungal species, of which *Cantharellus cibarius*, *Tricholoma magnivelare*, *Boletus edulis*

389 *s.l.* and *Lactarius deliciosus s.l.* are the most remarkable epigeous edible species, apart from the  
390 hypogeous fungi *Tuber melanosporum* and *Tuber aestivum*.

391 Some experimental plots with trees mycorrhized with *Cantharellus cibarius* were established in  
392 Sweden. The *C. cibarius* mycorrhiza survived transplantation and spread onto new roots but  
393 sporocarps did not form (Danell et al., 2002, Wang and Chen, 2014).

394 Plants mycorrhized with *Tricholoma matsutake* have been produced *in vitro*, and mycorrhizal  
395 formation *in situ* has also been reported (Wang et al., 1997; Yamada et al., 2006; Wang et al., 2012).  
396 More than 40,000 mycorrhized seedlings have been planted into existing *Pinus densiflora* forests in  
397 South Korea where *T. matsutake* is not currently present. However, there is no report of the progress  
398 of this experiment (Wang and Chen, 2014) and, to date, no other attempts to increase the production  
399 of this species in forests using this method have been reported.

400 With regard to *Boletus* species, mycorrhized plants have been produced under controlled conditions  
401 (Zuccherelli, 1988; Olivier et al., 1997). Some plantations have also been established in  
402 Mediterranean countries, such as Spain, where *Cistus* plants mycorrhized with *Boletus edulis*  
403 produced fruiting bodies three years after transplanting (Oria-de-Rueda et al., 2008), as well as in  
404 other small-scale experiments around the world (Wang and Chen, 2014). Recently, Mediavilla et al.  
405 (2016) observed that mycorrhization of *B. edulis* species could be improved using the helper bacteria  
406 *Pseudomonas fluorescens*.

407 The cultivation of *Lactarius* species has been widely studied since the 1980s because of their  
408 importance for recreational picking activities and as a commercial harvest. Poitou et al. (1984)  
409 reported the production of fruiting bodies in the field from outplanted *Pinus pinaster* mycorrhized  
410 seedlings. Guinberteau et al. (1990) reported that sporocarps develop 3 years after the seedlings  
411 establish. The *Lactarius deliciosus s.l.* species is also cultivated in French and Italian orchards, with  
412 up to 80 kg ha<sup>-1</sup> harvested in the first years after transplanting (<http://www.robinpepinieres.com>). In

413 New Zealand, hundreds of hectares have been planted with trees inoculated with *L. deliciosus* over  
414 the past 10 years that produce fruiting bodies every year (Wang and Chen, 2014). For example, yields  
415 of 50 kg ha<sup>-1</sup> of *L. deliciosus* in *P. pinaster* plantations have been reported (Wang and Hall, 2004),  
416 as well as yields of *Lactarius* spp. of approximately 300 kg ha<sup>-1</sup> in 3-year-old *Pinus radiata*  
417 plantations in New Zealand (Guerin-Laguet et al., 2014). However, even for *Boletus* and *Lactarius*  
418 species, positive results are reported only for orchards and plantations on bare soil. There is no  
419 scientific evidence that planting mycorrhized trees in a natural forest ecosystem enhances mushroom  
420 production.

421 Other measures aimed at increasing the abundance of mycorrhized trees in a forest involve directly  
422 inoculating the soil with spores or retaining mature sporocarps within the forest to act as a source of  
423 inoculum. These procedures have been used in Japan to increase the productivity of *T. matsutake*  
424 (Wang and Hall, 2004). Similar measures have been employed in South Korea and China (Wang and  
425 Hall, 2004). In China, e.g., the yield of *Lactarius volemus* sporocarps increased in areas with mature  
426 pine trees inoculated with spores of this species (Liu et al., 2007). However, there is not a convincing  
427 body of evidence that infecting trees with target fungi at the time of planting or directly in the field  
428 within the first months after planting leads to infection and colonization (Duñabeitia et al., 2004).  
429 Indeed, plant inoculation with edible mycorrhizal mushrooms is mostly conducted using *in vitro* and  
430 greenhouse techniques.

### 431 **3.4 Other management practices and disturbances**

432 In addition to the above-mentioned stand characteristics and silvicultural treatments influencing  
433 fungal fruiting and/or contributing to changes in mycocenoses, additional human interventions related  
434 to forest management may also affect productivity, i.e., timber harvesting operations, mushroom  
435 picking activities, trampling, disturbance of woody debris and fire (including both wildfire and  
436 prescribed burning).

#### 437 3.4.1. *Timber harvest and detrimental soil disturbance*

438 Timber harvesting is one of the operations that may have a great influence on mushroom occurrence  
439 immediately after thinning or regeneration cuts. It is well known that the physical properties of the  
440 soil can be significantly affected by harvesting operations, depending on the load of the machinery  
441 and the frequency of passes (Picchio et al., 2012). Soil compaction caused by logging techniques  
442 results in a critical reduction of water retention capacity, thus negatively influencing the long-term  
443 dynamics and fruiting of ectomycorrhizal fungi such as *Cantharellus* spp. (Amaranthus et al., 1996).  
444 Therefore, the potential impact on soil compaction should be considered when planning thinning or  
445 regeneration cuts. In this regard, although frequent and light forest interventions may be preferable  
446 to support mushroom yield (Pilz and Molina, 2002), frequent harvesting operations may also result  
447 in greater soil compaction than infrequent heavy felling (Amaranthus et al., 1996).

448 Mushroom-oriented silviculture should therefore implement harvesting methods and procedures with  
449 the aim of preserving forest soils. Indeed, the use of low-impact harvesting procedures may reduce  
450 the negative effects of logging, and mycosilvicultural interventions may even generate positive effects  
451 by increasing the productivity of some target mushroom species (Bonet et al., 2012). For instance,  
452 motor-manual felling and the extraction of whole trees by yarder would be preferable to using heavy  
453 machinery or winching and skidding methods because the impact on soil compaction is minimal. If  
454 this procedure is not applicable or convenient, appropriate training, the use of directional felling and  
455 careful supervision are necessary to minimize harvesting impacts on forest soils (Marchi et al., 2014).

#### 456 3.4.2. *Mushroom picking*

457 Because of the increasing harvesting pressure in many parts of the world (Boa, 2004), a widespread  
458 concern about overharvesting and possible damage to fungal resources has been raised (Egli et al.,  
459 2006). Legal restrictions on the harvesting of edible fungi have been introduced in many countries  
460 because there is a fear that sporocarp removal, which often occurs before spore dispersal, might

461 impair reproduction. Two experiments conducted in Oregon (Norvell, 1995) and Switzerland (Egli et  
462 al., 1990; Egli et al., 2006) found that sporocarp yield was not related to levels of mushroom  
463 harvesting. In a recent study, Parladé et al. (2017) also found that intensive picking did not  
464 significantly affect the mycelial biomass of *Boletus edulis* in the soil.

465 The Oregon Mycological Society Chanterelle Study evaluated the effect of mushroom harvesting,  
466 including the impact of different harvesting methods (i.e., either pulling up or cutting), and found no  
467 significant productivity decline in the first 10 years since the start of the study (Norvell, 1995). Egli  
468 et al. (2006) also investigated whether the mushroom harvesting technique affected mushroom  
469 production. They found that systematic harvesting did not reduce future yields of sporocarps. Similar  
470 results have also been reported for specific mushroom species such as *Tricholoma magnivelare*  
471 (Luoma et al., 2006) and *Morchella* spp. (Pilz et al., 2004; Larson et al., 2016).

472 However, although mushroom harvesting does not appear to affect fruitbody production *per se*,  
473 trampling caused by successive mushroom pickers throughout a given forest area may negatively  
474 affect fruitbody production. Sporocarp abundance in trampled areas can be 70% lower than in  
475 untrampled sites according to Egli and Ayer (1997) and Egli et al. (2006). However, there is no  
476 evidence that trampling affects the abundance of soil mycelia. The authors suggested that trampling  
477 possibly damages the sporocarp primordia (Egli and Ayer, 1997). Another hypothesis is that the  
478 reduction in yield may be linked to a decrease in the water retention capacity in the upper soil layers  
479 due to soil compaction. However, some fungal species (e.g., *Morchella* species) are often observed  
480 to fruit more abundantly on disturbed or compacted forest soils. Thus, mushroom pickers seeking  
481 morels tend to follow paths of logging equipment to search for concentrations of morels, which  
482 sometimes fruit on the footprints of previous mushroom harvesters (Pilz et al., 2006).

#### 483 3.4.3. Removal of forest litter

Another disturbance that has been evaluated with regard to its effect on wild mushroom production is raking, i.e., the removal of forest litter. Several authors, such as Baar and Kuyper (1993) and Baar and Ter Braak (1996), have reported that mushroom production by mycorrhizal species in *Pinus sylvestris* stands increases as a result of litter removal, especially two years after litter removal. By contrast, Salerni and Perini (2004) reported that manual removal of the litter layer negatively affected the fruiting process of *Boletus edulis* in silver fir forests in Central Italy. These contrasting results may be because the fungal response to raking is species specific; however, as yet the effects of this operation are not well known.

#### 3.4.4. Removal of understory vegetation and shrub control

The aim of shrub control is to increase (i) the survival and growth rate of mycorrhized trees by reducing competition for water and nutrients and (ii) the proliferation of mycelium as a consequence of improving the growth conditions of the host trees'.

The removal of understory vegetation may have an influence on mushroom production in forest ecosystems. An experiment conducted in central Italy (Tuscany) on Turkey oak (*Quercus cerris*) stands between 2000 and 2002 reported that the yield of *Boletus aereus* increased after removing the shrub layer growing under the oak canopy (Nocentini et al., 2004). The authors suggested that the increase in yield was a consequence of the ecology of the species, i.e., *B. aereus* being a termophilous species would benefit from the increased sunlight reaching the soil.

However, shrub vegetation itself can also be very productive in terms of mushrooms. For instance, *Cistus* scrublands can provide high yields of valuable, edible mushroom species, including *Boletus edulis* (Hernández-Rodríguez et al., 2015a, 2015b). In this regard, shrubs may play an important role in mushroom-oriented silviculture inasmuch as they may act as an accompanying species/reservoir, hosting fungal species that may inoculate new trees emerging after thinning or regeneration cuts. Another potential use of shrubs is the restoration of non-productive, degraded land where trees are

not able to grow in order to obtain the joint benefits of soil protection (e.g., to reduce further soil erosion and degradation) and the production of highly valuable non-wood forest products such as mushrooms (Hernández-Rodríguez et al., 2013).

#### 3.4.5. Prescribed burning

Prescribed burning is a management practice that aims to remove deadwood and other fuel biomass with the main objective of reducing the fire risk. In some cases, this practice can enhance the production of certain mushroom species that benefit from major disturbances such as wildfires. For example, Pilz et al. (2004) found that some putative species belonging to the genus *Morchella* (identified by molecular techniques among harvested black morels) fruited on burned areas. Winder and Keefer (2008) found that the average morel production after fire is about 6,500 morels ha<sup>-1</sup>. However, where the level of duff consumption reached 71%, morel abundance increased up to 16,827 morels ha<sup>-1</sup>.

In a more recent study, Larson et al. (2016) assessed post-fire morel abundance in a mixed coniferous forest and estimated an average yield of approximately 1,700 morels ha<sup>-1</sup> in the first year following a fire. Moreover, they also demonstrated that additional key factors controlling post-fire morel production vary at small spatial scales, resulting in the spatial patchiness of morel distribution. However, not all *Morchella* species are fire-prone (Pilz et al., 2004). While sporocarp formation of some *Morchella* species is influenced solely by temperature and precipitation during springtime (Mihail et al., 2007), other species require an additional forest disturbance, including fire (Pilz et al., 2004; Wurtz et al., 2005). Furthermore, Pilz et al. (2007) suggest that morels can behave either as saprotrophic or mycorrhizal fungi at different stages of their life cycle.

Martín-Pinto et al. (2006) assessed the effect of wildfire on mushroom production in scrublands (dominated by *Cistus ladanifer*) and *Pinus pinaster* stands in central Spain. They observed an increase in sporocarp yield of pyrophytic mycorrhizal species (e.g., *Leccinum corsicum*), whereas *Boletus*

532 *edulis* was not found in burned plots. More recently, Hernández-Rodríguez et al. (2015a) evaluated  
533 the effect of fire on fungal production in *C. ladanifer* scrublands by comparing the effects of total  
534 burning versus total clearing of the vegetation. In the first years after treatment, they found that  
535 mushroom production by mycorrhizal fungi was significantly higher after total clearing as compared  
536 to total burning. The authors suggest that if the primary objective is to increase the economic  
537 profitability of *C. ladanifer* scrublands, total clearing may be a preferable management option in order  
538 to enhance the yield of valuable, edible mushrooms.

#### 539 3.4.6. *Absence of forest management*

540 In the long term, if no management practices are performed, stands may gradually evolve into old-  
541 growth forests. Unmanaged old-growth forests differ from managed stands because the life cycle of  
542 trees is not interrupted and natural dynamics resulting after disturbances (e.g., parasite outbreaks,  
543 forest fires and heavy windstorms, gap creation due to insects, fungal decay and windstorms) are not  
544 controlled by forest management (Heilmann-Clausen et al., 2017). From a structural point of view,  
545 old-growth forests differ from mature managed forests in terms of the amount of living biomass and  
546 in the richness and abundance of dead wood habitats.

547 Very few studies have evaluated mushroom yields in old-growth forests. Smith et al. (2002) observed  
548 very low sporocarp yields in old-growth *Pseudotsuga menziesii* forest stands, especially when  
549 compared to managed stands. Furthermore, edible species such as *L. deliciosus* were absent from old-  
550 growth forests that were more than 400 years old (Smith et al., 2002), which may be because this  
551 early-seral mushroom species prefers younger and rather open stands (Martínez-Peña et al., 2012a).

## 552 4. Conclusions and future perspectives

### 553 4.1 Current mycosilvicultural knowledge

554 Effective mycosilviculture for the production of edible mushrooms in forests should carefully take  
555 into account the variables affecting mushroom yield. Although the weather conditions largely drive



556 mushroom yields (Alday et al., 2017a, 2017b), stand characteristics can also influence fungal  
557 resources. (i) Yields of mycorrhizal fungi are dependent on forest age, as more abundant sporocarps  
558 are generally observed in younger stands, which have a higher growth rate than trees in older stands.  
559 (ii) Canopy cover affects fungal yields: important edible and marketable species (*L. deliciosus*)  
560 benefit from stands with open canopies. However, in very open stands, productivity is lower than in  
561 less open stands. (iii) Stand density is one of the variables that influences mushroom production.  
562 Suitable stand basal areas for mushroom production depend on forest ecosystems and fungal species.  
563 In pine ecosystems, stand basal areas close to 20 m<sup>2</sup> ha<sup>-1</sup> and 40 m<sup>2</sup> ha<sup>-1</sup> are optimal for *Lactarius*  
564 *deliciosus* s.l. and *Boletus edulis* s.l., respectively. (iv) Tree composition influences fungal  
565 communities due to the preference of some ectomycorrhizal and saprotrophic fungi for specific host  
566 trees or litter, respectively.

567 Forest management practices affect mushroom occurrence and production both positively and  
568 negatively (Figure 2). The main findings from the reviewed literature can be summarized as follows:

569 (i) The effect of thinning on fungal production is the most studied silvicultural practice. However, the  
570 effect of thinning on mushroom yield may be different for different fungal species. Several authors  
571 have found that thinning, particularly in coniferous stands, has a positive effect on the sporocarp yield  
572 of important marketable species (*Lactarius* spp., *Boletus* spp.) if it creates low-medium density  
573 stands. By contrast, a negative effect of thinning has been reported for target species such as  
574 *Cantharellus* spp., and high-intensity thinning can also lead to a reduction in sporocarp yield.

575 (ii) At least in the short-term, clear-cutting negatively affects mushroom production, mainly of  
576 mycorrhizal fungi. If several mature trees are kept within the stand as retention trees, the negative  
577 effect of this treatment could be reduced. For this reason, shelterwood methods may be preferable to  
578 support mushroom production during the regeneration period in even-aged stands.

- 579 (iii) The use of mycorrhized seedlings seems to be a promising way of increasing fungal production,  
580 especially after clear-cutting. However, while many positive results have been reported for  
581 greenhouse or plantation experiments, there is no clear scientific evidence that successfully  
582 transplanting mycorrhized plants in natural forests enhances mushroom production.
- 583 (iv) Due to soil compaction, logging negatively affects mushroom occurrence, especially if timber  
584 harvesting procedures are repeated frequently on the same site. Low-impact harvesting methods  
585 coupled with careful supervision are needed to minimize the impact of these operations on mushroom  
586 communities.
- 587 (v) Harvesting sporocarps does not have a direct impact on fungal yield, unless the ground is  
588 excessively trampled by mushroom pickers, which can result in a decrease in sporocarp yield. Further  
589 research is needed on the effects of trampling in areas visited by huge numbers of mushroom pickers  
590 to ensure the long-term provision of mushroom-based ecosystem services.
- 591 (vi) The impact of litter removal remains unclear: some studies have reported negative effects on the  
592 productivity of important marketable species (*Boletus edulis*), while other authors have reported the  
593 opposite trend for mycorrhizal species.
- 594 (vii) Fire can reduce mushroom production even more than clear-cutting. Exceptions are pyrophytic  
595 species, including, e.g., some strains of *Morchella* spp.

596

597 (FIGURE 2 HERE)

598

## 599 4.2 Gaps in knowledge and research needs

600 In this review we have summarized our current understanding of mushroom production and examined  
601 the main silvicultural effects that influence productivity (Table 2). We have identified several gaps in  
602 knowledge that require further research, which can be summarized in the following points.

603

604 (TABLE 2 HERE)

605

606 (i) Stand age is an important factor that can affect the mushroom production of socially and  
607 economically important fungal species (Bonet et al., 2004). However, there is a considerable gap in  
608 our knowledge concerning the influence of tree age distribution within the stand. Most studies  
609 conducted to date have focused on even-aged forest stands and, therefore, the whether there are  
610 differences in the productivity levels of even-aged and uneven- or multi-aged stands of a given forest  
611 ecosystem remains unclear. Given that most trees in uneven-aged stands belong to the younger age  
612 classes, such stands might benefit from the joint production of early- and late-successional fungi  
613 simultaneously, but further research is required to test this and alternative hypotheses.

614 (ii) Several studies have investigated the effect of silvicultural practices and logging operations in the  
615 first years after the intervention, but the long-term effect has not been fully explored. Further research  
616 should be devoted to understanding the recovery time of mushroom resources after different  
617 silvicultural operations (e.g., thinning or regeneration cuts). Moreover, further research on low-  
618 impact harvesting procedures in relation to wild epigeous mushrooms is also needed.

619 (iii) Most of the studies conducted to date have focused on sporocarp yield; however, a more  
620 comprehensive picture could be obtained if the abundance of mycelium in the soil was also taken into  
621 account (Liu et al., 2016; Parladé et al., 2017). Indeed, the disappearance of some species after logging  
622 may not be definitive, but simply relate to the new microclimatic conditions at soil level.

623 (iv) Most research conducted to date has focused on a few coniferous forest ecosystems (mainly  
624 *Pinus*-dominated stands). Less is known about other coniferous ecosystems (e.g. *Abies*-, *Larix*- and  
625 *Pseudotsuga*-dominated stands), particularly broadleaved forests such as beech, chestnut or oak  
626 forests. Deciduous forests are very important for mushroom picking activities in many regions, such  
627 as in temperate ecosystems and Mediterranean climates.

628 (v) To our knowledge, to date, no studies have been devoted to assessing the influence of tree species  
629 diversity on mushroom production. In particular, further research is needed to evaluate differences in  
630 fungal yield under pure and mixed forest ecosystems.

631 (vi) A comparison of the mushroom yields obtained under different silvicultural regimes is needed to  
632 verify, e.g., whether even-aged management regimes are better than uneven-aged ones or if coppice  
633 is preferable to high forest in broadleaved stands. This information would help managers to determine  
634 whether maintaining coppices or converting to high forest is the best option for mushroom yields and  
635 wood production.

636 (vii) The use of mycorrhized shrubs to restore degraded lands where trees are not able to grow may  
637 have promise as a way of increasing the value of degraded lands. However, to date, only a few studies  
638 have been carried out and further research is needed to provide further scientific evidence for different  
639 scrubland ecosystems.

640 (viii) Old-growth forests are considered to be an important resource that needs to be preserved,  
641 especially their role in biodiversity conservation. However, little is known about mushroom yields in  
642 old growth forests.

643 (ix) Most mycosilvicultural research studies have been conducted in Europe and North America,  
644 focusing on Mediterranean, temperate and boreal ecosystems. Our scientific understanding of  
645 mycosilviculture in other regions of the world and in other forest ecosystems and biomes (e.g.,  
646 tropical) is still very limited.

647 (x) Models to predict mushroom yields have been proposed for relevant marketable species (e.g.,  
648 *Boletus* spp. and *Lactarius* spp.); however, their validity is still local or regional. Silvicultural models  
649 that correlate arboreal carbon allocation with stand conditions and management regimes are available  
650 (e.g., Landsberg et al., 2003) and may be applied to model the production of forest mushrooms in  
651 order to provide more dynamic and efficient estimates of mushroom yields.

652 (xi) The context of global change should also be considered in future research. In particular, we need  
653 to determine how changes in precipitation and temperature patterns associated to climate change as  
654 well as global perturbations in the C and N cycles (increased CO<sub>2</sub> in the atmosphere; nitrogen  
655 deposition) will affect mushroom fruiting (Pickles et al., 2012; Suz et al., 2014). Furthermore, the  
656 predicted vegetation shifts, which may result in changes (i.e., migration) of host tree species of  
657 valuable fungi, should be also taken into consideration when modelling future mushroom yield  
658 scenarios.

659 In summary, a better understanding of the effect of forest management practices on mushroom  
660 production should provide a basis for improving productivity in forest stands. The research lines  
661 outlined above should not only enable us to acquire an in-depth ecological knowledge of forest  
662 ecosystems and fungal dynamics but also provide valuable scientific knowledge to support decision-  
663 making in forest management planning and policy-making (Corona, 2014; Corona et al., 2016).

## 664 **Acknowledgements**

665 The review was carried out within the COST Action FP1203 (European Non-Wood Forest Products).  
666 Sergio de-Miguel was supported by the European Union's Horizon 2020 MultiFUNGtionality Marie  
667 Skłodowska-Curie (IF-EF No-655815), by the Spanish research project AGL2015-66001-C3-1-R  
668 (MEC Spain) and by the European project StarTree (No. 311919). José Antonio Bonet benefited from  
669 a Serra-Hunter Fellowship provided by the Generalitat of Catalunya.

## 670 **References**

671 Ágreda, T., Cisneros, Ó., Águeda, B., Fernández-Toirán, L. M., 2014. Age class influence on the  
 672 yield of edible fungi in a managed Mediterranean forest. *Mycorrhiza*. 24(2), 143-152.

673 Ágreda, T., Águeda, B., Fernández-Toirán, M., Vicente-Serrano, S. M., Òlano, J. M., 2016. Long-  
 674 term monitoring reveals a highly structured interspecific variability in climatic control of sporocarp  
 675 production. *Agric. For. Meteor.* 223, 39-47.

676 Alday, J.G., Martínez de Aragón, J., de-Miguel, S., Bonet, J.A., 2017a. Mushroom biomass and  
 677 diversity are driven by different spatio-temporal scales along Mediterranean elevation gradients.  
 678 *Scientific Reports*, 7, 45824; doi: 10.1038/srep45824.

679 Alday, J.G., Bonet, J.A., Oria-de-Rueda, J.A., Martínez de Aragón, J., Aldea, J., Martín-Pinto, P., de-  
 680 Miguel, S., Hernández-Rodríguez, M., Martínez-Peña, F. 2017b. Record breaking mushroom yields  
 681 in Spain. *Fungal Ecol.* 26, 144-146; doi: 10.1016/j.funeco.2017.01.004.

682 Amaranthus, M. P., Page-Dumroese, D., Harvey, A., Cazares, E., Bednar, L. F., 1996. Soil  
 683 compaction and organic matter affect conifer seedling nonmycorrhizal and ectomycorrhizal root tip  
 684 abundance and diversity. Research Paper PNW-RP-494. U.S. Department of Agriculture, Forest  
 685 Service, Pacific Northwest Research Station, Portland, OR, p. 12.

686 Baar, J., Kuyper, T. W., 1993. Litter removal in forests and effect on mycorrhizal fungi. In *Fungi of*  
 687 *Europe: Investigation, Recording and Conservation* (eds. D. N. Pegler, L. Boddy, B. Ing and P. M.  
 688 Kirk), pp. 275–286. Royal Botanic Gardens, Kew.

689 Baar, J., Ter Braak, C. J. F., 1996. Ectomycorrhizal sporocarp occurrence as affected by manipulation  
 690 of litter and humus layers in Scots pine stands of different age. *Appl. Soil Ecol.* 4, 61–73.

691 Barluzzi, C., Perini, C., De Dominicis, V., 1992. Coenological research on macrofungi in chestnut  
 692 coppices of Tuscany. *Phytocoenologia*, 20(4), 449-465.

693 Barroetaveña, C., La Manna, L., Alonso, M. V., 2008. Variables affecting *Suillus luteus* fructification  
694 in ponderosa pine plantations of Patagonia (Argentina). *For. Ecol. Manage.* 256(11), 1868-1874.

695 Bengtsson, J., Nilsson, S. G., Franc, A., Menozzi, P., 2000. Biodiversity, disturbances, ecosystem  
696 function and management of European forests. *For. Ecol. Manage.* 132(1), 39-50.

697 Bettini, G., Bianchetto, E., Butti, F., Chiellini, C., De Meo, I., d'Errico, G., Fabiani, A., Gardin, L.,  
698 Graziani, A., Landi, S., Marchi, M., Mazza, G., Mocali, S., Montini, P., Plutino, M., Roversi, P. F.,  
699 Salerni, E., Samaden, S., Canencia, I. S., Torrini, G., 2016. Selective thinning: Increasing mechanical  
700 stability and biodiversity in black pine plantations. Ed. Cantiani P., Compagnia delle Foreste Srl.  
701 Available on line at [[http://www.selpibio.eu/en/publications/item/43-technical-handbook-selective-](http://www.selpibio.eu/en/publications/item/43-technical-handbook-selective-thinning.html)  
702 [thinning.html](http://www.selpibio.eu/en/publications/item/43-technical-handbook-selective-thinning.html)]

703 Boa, E. R., 2004. Wild edible fungi: a global overview of their use and importance to people (No.  
704 17). Food & Agriculture Org.

705 Bonet, J. A., Fischer, C. R., Colinas, C., 2004. The relationship between forest age and aspect on the  
706 production of sporocarps of ectomycorrhizal fungi in *Pinus sylvestris* forests of the central Pyrenees.  
707 *For. Ecol. Manage.* 203(1), 157-175.

708 Bonet, J. A., Pukkala, T., Fischer, C. R., Palahí, M., Martínez de Aragón, J., Colinas, C., 2008.  
709 Empirical models for predicting the production of wild mushrooms in Scots pine (*Pinus sylvestris* L.)  
710 forests in the Central Pyrenees. *Ann. For. Sci.* 65(2), 206-206.

711 Bonet, J. A., Oliach, D., Fischer, C., Olivera, A., Martínez de Aragón, J., Colinas, C., 2009.  
712 Cultivation methods of the black truffle, the most profitable Mediterranean non-wood forest product;  
713 a state of the art review. *EFI Proc.* 57, 57-71.

714 Bonet, J. A., Palahí, M., Colinas, C., Pukkala, T., Fischer, C. R., Miina, J., Martínez de Aragón, J.,  
 715 2010. Modelling the production and species richness of wild mushrooms in pine forests of the Central  
 716 Pyrenees in northeastern Spain. *Can. J. For. Res.*, 40(2), 347-356.

717 Bonet, J. A., De-Miguel, S., de Aragón, J. M., Pukkala, T., Palahí, M., 2012. Immediate effect of  
 718 thinning on the yield of *Lactarius group deliciosus* in *Pinus pinaster* forests in Northeastern Spain.  
 719 *For. Ecol. Manage.*, 265, 211-217.

720 Bonet, J. A., González-Olabarria, J. R., Martínez de Aragón, J., 2014. Mushroom production as an  
 721 alternative for rural development in a forested mountainous area. *J. Mt. Sci.* 11(2), 535-543.

722 Buée, M., Maurice, J. P., Marçais, B., Dupouey, J. L., Garbaye, J., Le Tacon, F., 2005. Effet des  
 723 interventions sylvicoles sur les champignons sylvestres. *Forêt-entreprise*. 164, 26-32.

724 Buée, M., Maurice, J.P., Zeller, B., Andrianarisoa, S., Ranger, J., Courtecuisse, R., Marçais, B., Le  
 725 Tacon, F., 2011. Influence of tree species on richness and diversity of epigeous fungal communities  
 726 in a French temperate forest stand. *Fungal Ecol.* 4, 22–31.

727 Büntgen, U., Kauserud, H., Egli, S., 2012. Linking climate variability to mushroom productivity and  
 728 phenology. *Front. Ecol. Environ.* 10, 14–19.

729 Cai, M., Pettenella, D., Vidale, E., 2011. Income generation from wild mushrooms in marginal rural  
 730 areas. *For. Policy Econ.* 13, 211–226.

731 Chibisov, G., Demidova, N., 1998. Non-wood forest products and their research in Arkhangelsk,  
 732 Russia. *EFI Proc.* 23, 147-153.

733 Ciancio, O. 1981. Le specie forestali esotiche e le relazioni tra arboricoltura e selvicoltura. *Ann. Ist.*  
 734 *Sper. Selvic.* 12-13: 1-100. (in Italian)



735 Ciancio, O., Nocentini, S., 1997. The forest and man: the evolution of forestry thought from modern  
 736 humanism to the culture of complexity. Systemic silviculture and management on natural bases. The  
 737 forest and man. Florence, Accademia Italiana di Scienze Forestali, 21-114.

738 Corona, P., 2015. Forestry research to support the transition towards a bio-based economy. Ann.  
 739 Silv. Res. 38(2), 37-38.

740 Corona, P., Cutini, A., Chiavetta, U., Paoletti, E., 2016. Forest-food nexus: a topical opportunity for  
 741 human well-being and silviculture. Ann. Silv. Res. 40(1), 1-10.

742 Courtecuisse, R., Duhem, B., 1994. Guide des champignons de France et d'Europe. Lausanne:  
 743 Delachaux et Niestlé.

744 Danell, E., Hall, I., Wang, Y., Zambonelli, A., 2002. Current research on chanterelle cultivation in  
 745 Sweden. In Edible mycorrhizal mushrooms and their cultivation. Proceedings of the Second  
 746 International Conference on Edible Mycorrhizal Mushrooms, Christchurch, New Zealand, 3-6 July,  
 747 2001. pp.: 0-4. Crop & Food Res.

748 de Groot, M., Eler, K., Flajšman, K., Grebenc, T., Marinšek, A., Kutnar, L., 2016. Differential short-  
 749 term response of functional groups to a change in forest management in a temperate forest. For. Ecol.  
 750 Manage. 376, 256-264.

751 de-Miguel, S., Bonet, J. A., Pukkala, T., Martínez de Aragón, J., 2014. Impact of forest management  
 752 intensity on landscape-level mushroom productivity: a regional model-based scenario analysis. For.  
 753 Ecol. Manage. 330, 218-227.

754 De Román, M., Boa, E., 2006. The marketing of *Lactarius deliciosus* in northern Spain. Econ. Bot.  
 755 60(3), 284-290.

756 Dettori, S., Marone, E., Portoghesi, L., 2009. Filiera delle produzioni forestali non legnose:  
 757 produzione e raccolta tra sostenibilità e tracciabilità, in: Proceedings, Terzo Congresso Nazionale Di

758 Selvicoltura. Accademia Italiana Di Scienze Forestali, Ministero Delle Politiche Agricole, Alimentari  
 759 E Forestali, Corpo Forestale Dello Stato, Ministero dell’Ambiente, Della Tutela Del Territorio E Del  
 760 Mare, Regi. pp. 742–751.

761 Dighton, J., Mason, P. A., 1985. Mycorrhizal dynamics during forest tree development. In: Moore D,  
 762 Castellano LA, Wood D, Frankland JC (eds) Development biology of higher fungi. University Press  
 763 Cambridge, Cambridge, pp 117–139.

764 Duñabeitia, M., Rodríguez, N., Salcedo, I., Sarrionandia, E., 2004. Field mycorrhization and its  
 765 influence on the establishment and development of the seedlings in a broadleaf plantation in the  
 766 Basque Country. *For. Ecol. Manage.*, 195(1), 129-139.

767 Durall, D.M., Gamiet, S., Simard, S.W., Kudrna, L., Sakakibara, S.M., 2006. Effects of clear-cut  
 768 logging and tree species composition on the diversity and community composition of epigeous fruit  
 769 bodies formed by ectomycorrhizal fungi. *Can. J. Bot.* 84, 966–980.

770 Duryea, M. L., Dougherty, P. M., 2012. Forest regeneration manual (Vol. 36). Springer Science &  
 771 Business Media.

772 Egli, S., Ayer, F., Chatelain, F., 1990. Der einfluss des pilzsammelns auf die pilzflora. *Mycol. Helv.*  
 773 3(4), 417-428. (In German).

774 Egli, S., Ayer, F., 1997. Est-il possible d'améliorer la production de champignons comestibles en  
 775 forêt? L'exemple de la réserve mycologique de La Chanéaz en Suisse. *Rev. For. Fr.* 49, 235-243.

776 Egli, S., Peter, M., Buser, C., Stahel, W., Ayer, F., 2006. Mushroom picking does not impair future  
 777 harvests—results of a long-term study in Switzerland. *Biol. Conserv.* 129(2), 271-276.

778 Egli, S., Ayer, F., Peter, M., Eilmann, B., Rigling, A., 2010. Is forest mushroom productivity driven  
 779 by tree growth? Results from a thinning experiment. *Ann. For. Sci.*, 67(5), 509.

780 Egli, S., 2011. Mycorrhizal mushroom diversity and productivity—an indicator of forest health? Ann.  
781 For. Sci. 68(1), 81-88.

782 Fernández-Toirán, L. M., Ágreda, T., Olano, J. M., 2006. Stand age and sampling year effect on the  
783 fungal fruit body community in *Pinus pinaster* forests in central Spain. Bot. 84(8), 1249-1258.

784 Gange, A. C., Gange, E. G., Sparks, T. H., Boddy, L., 2007. Rapid and recent changes in fungal  
785 fruiting patterns. Sci. 316(5821), 71-71.

786 Giovannetti, G., Roth-Bejerano, N., Zanini, E., Kagan-Zur, V. (2010). Truffles and their cultivation.  
787 Hortic. Rev. 16, 71-107.

788 Grebenc, T., Christensen, M., Vilhar, U., Čater, M., Martín, M. P., Simoncic, P., Kraigher, H., 2009.  
789 Response of ectomycorrhizal community structure to gap opening in natural and managed temperate  
790 beech-dominated forests. Can. J. For. Res. 39(7), 1375-1386.

791 Guerin-Laguette, A., Cummings, N., Butler, R. C., Willows, A., Hesom-Williams, N., Li, S., Wang,  
792 Y., 2014. *Lactarius deliciosus* and *Pinus radiata* in New Zealand: towards the development of  
793 innovative gourmet mushroom orchards. Mycorrhiza. 24(7), 511-523.

794 Guinberteau, J., Ducamp, M., Poitou, N., Mamoun, M., Olivier, J. M., 1990. Ecology of various  
795 competitors from an experimental plot of *Pinus pinaster* inoculated with *Suillus granulatus* and  
796 *Lactarius deliciosus*. Agriculture, Ecosyst. Environ. 28(1), 161-165.

797 Harvey, A. E., Jurgensen, M. F., Larsen, M. J., 1980. Clearcut harvesting and ectomycorrhizae:  
798 survival of activity on residual roots and influence on a bordering forest stand in western Montana.  
799 Can. J. For. Res. 10(3), 300-303.

800 Heilmann-Clausen, J., Adamčík, S., Bässler, C., Halme, P., Krisai-Greilhuber, I., Holec, J., 2017.  
801 State of the art and future directions for mycological research in old-growth forests. Fungal Ecol. 27,  
802 141-144.

803 Hernández-Rodríguez, M., Oria-de-Rueda, J. A., Martín-Pinto, P., 2013. Post-fire fungal succession  
804 in a Mediterranean ecosystem dominated by *Cistus ladanifer* L. For. Ecol. Manage. 289, 48-57.

805 Hernández-Rodríguez, M., de Miguel, S., Pukkala, T., Oria de Rueda, J.A., Martín-Pinto, P., 2015a.  
806 Climate-sensitive models for mushroom yields and diversity in *Cistus ladanifer* scrublands. Agric.  
807 For. Meteorol. 213, 173–182.

808 Hernández-Rodríguez, M., Oria-de-Rueda, J. A., Pando, V., Martín-Pinto, P., 2015b. Impact of fuel  
809 reduction treatments on fungal sporocarp production and diversity associated with *Cistus ladanifer*  
810 L. ecosystems. For. Ecol. Manage. 353, 10-20.

811 Hintikka, V., 1988. On the macromycete flora in oligotrophic pine forests of different ages in South  
812 Finland. Acta Bot. Fennica 136, 89–94.

813 ISTAT, 2008. Bollettino mensile di statistica, n. 3 marzo 2008, Istituto Nazionale di Statistica, Rome.  
814 Retrieved from:  
815 [http://www.istat.it/dati/catalogo/20080807\\_02/Bollettino%20istat%20marzo%2008.pdf](http://www.istat.it/dati/catalogo/20080807_02/Bollettino%20istat%20marzo%2008.pdf) (in italian).

816 Kalamees, K., Silver, S., 1988. Fungal productivity of pine heaths in North-West Estonia. Acta Bot.  
817 Fennica 136, 95–98.

818 Kardell, L., Eriksson, L., 1987. Kremlor, riskor, soppar. Skogsbruksmetodernas inverkan på  
819 produktionen av matsvampar. Sveriges Skogsvårdsförbunds Fidskrift, 2(87), 3-24. (In Swedish)

820 Kauserud, H., Stige, L.C., Vik, J.O., Okland, R.H., Hoiland, K., Stenseth, N.C., 2008. Mushroom  
821 fruiting and climate change. Proc. Natl. Acad. Sci. U.S.A. 105, 3811–3814.

822 Keizer, P. J., Arnolds, E., 1994. Succession of ectomycorrhizal fungi in roadside verges planted with  
823 common oak (*Quercus robur* L.) in Drenthe, The Netherlands. Mycorrhiza, 4(4), 147-159.

824 Kennedy, P. G., Izzo, A. D., Bruns, T. D., 2003. There is high potential for the formation of common  
 825 mycorrhizal networks between understorey and canopy trees in a mixed evergreen forest. J. Ecol.  
 826 91(6), 1071-1080.

827 Kim, M.S., Klopfenstein, N.B., McDonald, G.I., 2010. Effects of forest management practices and  
 828 environment on occurrence of Armillaria Species. Jour. Korean For. Soc. 99 (2), 251-257.

829 Kranabetter, J. M., Trowbridge, R., Macadam, A., McLennan, D., Friesen, J. 2002. Ecological  
 830 descriptions of pine mushroom (*Tricholoma magnivelare*) habitat and estimates of its extent in  
 831 northwestern British Columbia. For. Ecol. Manage. 158(1), 249-261.

832 Kranabetter, J. M., Friesen, J., Gamiet, S., Kroeger, P., 2005. Ectomycorrhizal mushroom distribution  
 833 by stand age in western hemlock-lodgepole pine forests of northwestern British Columbia. Can. J.  
 834 For. Res. 35(7), 1527-1539.

835 Kranabetter, J. M., Williams, H., Morin, J., 2009. Ecological descriptions of Pacific golden  
 836 chanterelle (*Cantharellus formosus*) habitat and estimates of its extent in Haida Gwaii. J. Ecosyst.  
 837 Manag. 10(1), 59–67.

838 Krebs, C. J., Carrier, P., Boutin, S., Boonstra, R., Hofer, E., 2008. Mushroom crops in relation to  
 839 weather in the southwestern Yukon. Bot. 86(12), 1497-1502.

840 Kropp, B. R., Albee, S., 1996. The effects of silvicultural treatments on occurrence of mycorrhizal  
 841 sporocarps in a *Pinus contorta* forest: a preliminary study. Biol. Conserv. 78(3), 313-318.

842 Kucuker, D. M., Baskent, E. Z., 2015. Spatial prediction of *Lactarius deliciosus* and *Lactarius*  
 843 *salmonicolor* mushroom distribution with logistic regression models in the Kızılıcasu Planning Unit,  
 844 Turkey. Mycorrhiza. 25(1), 1-11.

845 Landi, M., Salerni, E., Ambrosio, E., D'Aguanno, M., Nucci, A., Saveri, C., Perini, C., Angiolini, C.,  
 846 2015. Concordance between vascular plant and macrofungal community composition in broadleaf  
 847 deciduous forests in central Italy. *iForest-Biogeosciences and For.* 8(3), 279.

848 Landsberg, J. J., Waring, R. H., Coops, N. C., 2003. Performance of the forest productivity model 3-  
 849 PG applied to a wide range of forest types. *For. Ecol. Manage.* 172(2), 199-214.

850 Larson, A. J., Cansler, C. A., Cowdery, S. G., Hiebert, S., Furniss, T. J., Swanson, M. E., Lutz, J. A.,  
 851 2016. Post-fire morel (*Morchella*) mushroom abundance, spatial structure, and harvest sustainability.  
 852 *For. Ecol. Manage.* 377, 16-25.

853 Latorre, J., 2016. Mycotourism in Castilla y León – an innovative opportunity to stimulate the  
 854 development of rural areas. NWFP Update, 10 (on-line version).

855 Lee, C. A., Dey, D. C., Muzika, R. M., 2016. Oak stump-sprout vigor and *Armillaria* infection after  
 856 clearcutting in southeastern Missouri, USA. *For. Ecol. Manage.* 374, 211-219.

857 Le Tacon, F., Alvarez, I.F., Bouchard, D., Henrion, B., Jackson, R.M., Luff, S., Parlade, J., Pera, J.,  
 858 Sternström, E., Villeneuve, N., Walker, C., 1992. Variations in field responses of forest trees to  
 859 nursery ectomycorrhizal inoculation in Europe. In: Read DJ, Lewis DH, Lewis AH, Alexander IJ  
 860 (eds) *Mycorrhizas in ecosystems*. CAB International, Wallingford, pp: 119–134.

861 Le Tacon, F., 1997. Vers une meilleure prise en compte des champignons mycorhiziens dans la  
 862 gestion forestière. *Rev. For. Fr.* 49, 245-255.

863 Lin, W. R., Wang, P. H., Chen, M. C., Kuo, Y. L., Chiang, P. N., Wang, M. K., 2015. The impacts  
 864 of thinning on the fruiting of saprophytic fungi in *Cryptomeria japonica* plantations in central Taiwan.  
 865 *For. Ecol. Manage.* 336, 183-193.

866 Liu, B., Bonet, J. A., Fischer, C. R., de Aragón, J. M., Bassie, L., Colinas, C., 2016. *Lactarius*  
 867 *deliciosus* Fr. soil extraradical mycelium correlates with stand fruitbody productivity and is increased  
 868 by forest thinning. For. Ecol. Manage. 380, 196-201.

869 Liu, P.G., Yu, F.Q., Wang, X.H., Chen, J., Zheng, H.D., Tian, X.F., 2007. Mycorrhizal edible fungi  
 870 stimulate solutions and their preparation. China Patents, Patent No. CN1958784A

871 Luoma, D. L., Eberhart, J. L., Molina, R., Amaranthus, M. P., 2004. Response of ectomycorrhizal  
 872 fungus sporocarp production to varying levels and patterns of green-tree retention. For. Ecol. Manage.  
 873 202(1), 337-354.

874 Luoma, D. L., Eberhart, J. L., Abbott, R., Moore, A., Amaranthus, M. P., Pilz, D., 2006. Effects of  
 875 mushroom harvest technique on subsequent American matsutake production. For. Ecol. Manage.  
 876 236(1), 65-75.

877 Mäkinen, H., Isomäki, A., 2004. Thinning intensity and growth of Scots pine stands in Finland. For.  
 878 Ecol. Manage. 201(2), 311-325.

879 Marchi, E., Picchio, R., Spinelli, R., Verani, S., Venanzi, R., Certini, G., 2014. Environmental impact  
 880 assessment of different logging methods in pine forests thinning. Ecol. Eng. 70, 429-436.

881 Martínez de Aragón, J., Bonet, J. A., Fischer, C. R., Colinas, C., 2007. Productivity of  
 882 ectomycorrhizal and selected edible saprotrophic fungi in pine forests of the pre-Pyrenees mountains,  
 883 Spain: predictive equations for forest management of mycological resources. For. Ecol. Manage.  
 884 252(1), 239-256.

885 Martínez de Aragón, J., Riera, P., Giergiczny, M., Colinas, C., 2011. Value of wild mushroom picking  
 886 as an environmental service. For. Policy Econ. 13(6), 419-424.

887 Martínez de Aragón, J., Fischer, C., Bonet, J. A., Olivera, A., Oliach, D., Colinas, C., 2012.  
 888 Economically profitable post fire restoration with black truffle (*Tuber melanosporum*) producing  
 889 plantations. New For. 43(5-6), 615-630.

890 Martínez-Peña, F., Oria de Rueda, J.A., Ágreda, T. (coords). 2011. Manual para la gestión del recurso  
 891 micológico forestal en Castilla y León. Serie Técnica de la Junta de Castilla y León. Valladolid. 453  
 892 pp.

893 Martínez-Peña, F., Ágreda, T., Águeda, B., Ortega-Martínez, P., Fernández-Toirán, L. M., 2012a.  
 894 Edible sporocarp production by age class in a Scots pine stand in Northern Spain. Mycorrhiza. 22(3),  
 895 167-174.

896 Martínez-Peña, F., De-Miguel, S., Pukkala, T., Bonet, J. A., Ortega-Martínez, P., Aldea, J., Martínez  
 897 de Aragón, J. M., 2012b. Yield models for ectomycorrhizal mushrooms in *Pinus sylvestris* forests  
 898 with special focus on *Boletus edulis* and *Lactarius group deliciosus*. For. Ecol. Manage. 282, 63-69.

899 Martín-Pinto, P., Vaquerizo, H., Peñalver, F., Olaizola, J., Oria-de-Rueda, J. A., 2006. Early effects  
 900 of a wildfire on the diversity and production of fungal communities in Mediterranean vegetation types  
 901 dominated by *Cistus ladanifer* and *Pinus pinaster* in Spain. For. Ecol. Manage. 225(1), 296-305.

902 Maso, D., 2008. Networks and PES schemes as income-generation tools for the development of the  
 903 Italian forest sector. PhD thesis. Retrieved from  
 904 [http://paduaresearch.cab.unipd.it/1377/1/Tesi\\_Daria\\_Maso.pdf](http://paduaresearch.cab.unipd.it/1377/1/Tesi_Daria_Maso.pdf).

905 Mason, P., Last, F., Wilson, J., 1987. Ectomycorrhizal fungi. In Fungal Infection of Plants:  
 906 Symposium of the British Mycological Society. Cambridge University Press. p. 253.

907 Mediavilla, O., Olaizola, J., Santos-del-Blanco, L., Oria-de-Rueda, J. A., Martín-Pinto, P., 2016.  
 908 Mycorrhization between *Cistus ladanifer* L. and *Boletus edulis* Bull is enhanced by the mycorrhiza  
 909 helper bacteria *Pseudomonas fluorescens* Migula. Mycorrhiza. 26(2), 161-168.



910 Mihail, J.D., Bruhn, J.N., Bonello, P., 2007. Spatial and temporal patterns of morel fruiting. Mycol.  
 911 Res. 111, 339–346.

912 Molina, R. 1979. Pure culture synthesis and host specificity of red alder mycorrhizae. Can. J Bot.  
 913 57(11), 1223-1228.

914 Molina, R., Trappe, J. M., 1982. Patterns of ectomycorrhizal host specificity and potential among  
 915 Pacific Northwest conifers and fungi. For. Sci. 28(3), 423-458.

916 Nocentini, G., Di Cocco, S., Di Cocco, G., 2004. Increasing the production of *Boletus aereus* in a  
 917 deciduous forest. The experience in the area of Mondeggi (FI). Sherwood. 10(11): 33-38 (in Italian)

918 Norvell, L., 1995. Loving the chanterelle to death? The ten-year Oregon chanterelle project.  
 919 McIlvainea, 12(1), 6-25.

920 Ogaya, R., Peñuelas, J., 2005. Decreased mushroom production in a holm oak forest in response to  
 921 an experimental drought. For. 78(3), 279-283.

922 Ohenoja, E., 1988. Effect of forest management procedures on fungal fruit body production in  
 923 Finland. Acta Bot. Fennica. 136, 81-84.

924 Olivera, A., Bonet, J. A., Oliach, D., Colinas, C., 2014. Time and dose of irrigation impact *Tuber*  
 925 *melanosporum* ectomycorrhiza proliferation and growth of *Quercus ilex* seedling hosts in young  
 926 black truffle orchards. Mycorrhiza, 24(1), 73-78.

927 Olivier, J. M., Guinberteau, J., Rondet, J., Mamoun, M., 1997. Vers l'inoculation contrôlée des cèpes  
 928 et bolets comestibles? Rev. For. Fr. 49, 222-234.

929 Oria-de-Rueda, J. A., Martín-Pinto, P., Olaizola, J., 2008. Bolete Productivity of Cistaceous  
 930 Scrublands in Northwestern Spain. Econ. Bot. 62(3), 323-330.

931 Ortega-Martínez, P., Águeda, B., Fernández-Toirán, L. M., Martínez-Peña, F., 2011. Tree age  
 932 influences on the development of edible ectomycorrhizal fungi sporocarps in *Pinus sylvestris* stands.  
 933 Mycorrhiza, 21(1), 65-70.

934 Packham, J. M., May, T. W., Brown, M. J., Wardlaw, T. J., Mills, A. K., 2002. Macrofungal diversity  
 935 and community ecology in mature and regrowth wet eucalypt forest in Tasmania: a multivariate study.  
 936 Austral Ecol. 27(2), 149-161.

937 Palahí, M., Pukkala, T., Bonet, J. A., Colinas, C., Fischer, C. R., Martínez de Aragón, J. R., 2009.  
 938 Effect of the inclusion of mushroom values on the optimal management of even-aged pine stands of  
 939 Catalonia. For. Sci. 55(6), 503-511.

940 Parladé, J., Martínez-Peña, F., Pera, J., 2017. Effects of forest management and climatic variables on  
 941 the mycelium dynamics and sporocarp production of the ectomycorrhizal fungus *Boletus edulis*. For.  
 942 Ecol. Manage. 390, 73-79.

943 Peter, M., Buée, M., Egli, S., 2013. Biodiversity of mycorrhizal fungi as a crucial player in forest  
 944 ecosystem functioning. In: Kraus, D., Krumm, F. (Eds.), Integrative approaches as an opportunity for  
 945 the conservation of forest biodiversity. Eur. For. Inst. pp. 170–179.

946 Pettenella, D., Secco, L., 2006. Small-scale forestry in the Italian Alps: from mass market to territorial  
 947 marketing. In: Wall, S. (Ed.), Small-scale forestry and rural development. The intersection of  
 948 ecosystems, economics and society, Proceedings of IUFRO 3.08 Conference, Galway, Galway-Mayo  
 949 Institute of Technology. Galway-Mayo Institute of Technology, Galway, pp. 398–408.

950 Picchio, R., Neri, F., Petrini, E., Verani, S., Marchi, E., Certini, G., 2012. Machinery-induced soil  
 951 compaction in thinning two pine stands in central Italy. For. Ecol. Manage. 285, 38-43.

952 Pickles, B. J., Egger, K. N., Massicotte, H. B., Green, D. S. 2012. Ectomycorrhizas and climate  
 953 change. Fungal Ecol. 5(1), 73-84.

954 Pierangelo, A., Rolland, B. 2013. Guida pratica di micoselvicultura. Available at:  
 955 [www.amycoforest.eu](http://www.amycoforest.eu) (In Italian)

956 Pilz, D., Molina, R., 2002. Commercial harvests of edible mushrooms from the forests of the Pacific  
 957 Northwest United States: issues, management, and monitoring for sustainability. *For. Ecol. Manage.*  
 958 155(1), 3-16.

959 Pilz, D., Weber, N. S., Carter, M. C., Parks, C. G., Molina, R., 2004. Productivity and diversity of  
 960 morel mushrooms in healthy, burned, and insect-damaged forests of northeastern Oregon. *For. Ecol.*  
 961 *Manage.* 198(1), 367-386.

962 Pilz, D., Molina, R., Mayo, J., 2006. Effects of thinning young forests on chanterelle mushroom  
 963 production. *J. For.* 104(1), 9-14.

964 Pilz, D., McLain, R., Alexander, S., Villarreal-Ruiz, L., Berch, S., Wurtz, T.L., Parks, C.G.,  
 965 McFarlane, E., Baker, B., Molina, R., Smith, J.E., 2007. Ecology and Management of Morels  
 966 Harvested from the Forests of western North America. USDA Forest Service General Technical  
 967 Report PNW-GTR-710.

968 Poitou, N., Mamoun, M., Ducamp, M., Delmas, J., 1984. After *Boletus granulatus*, *Lactarius*  
 969 *deliciosus* obtained in fructification in the field from mycorrhizal plants. *Rev. Hortic.* 244:65–68. (In  
 970 French)

971 Primicia, I., Camarero, J. J., Martínez de Aragón, J., de-Miguel, S., Bonet, J. A., 2016. Linkages  
 972 between climate, seasonal wood formation and mycorrhizal mushroom yields. *Agric. For. Meteorol.*  
 973 228, 339-348.

974 Pukkala, T., Von Gadow, K., 2012. Continuous Cover Forestry. Springer, Dordrecht.

975 Reyna, S., Garcia-Barreda, S., 2014. Black truffle cultivation: a global reality. *For. Syst.* 23(2), 317-  
 976 328.

977 Salerni, E., Perini, C., 2004. Experimental study for increasing productivity of *Boletus edulis* s.l. in  
978 Italy. For. Ecol. Manage. 201(2), 161-170.

979 Santos-Silva, C., Gonçalves, A., Louro, R., 2011. Canopy cover influence on macrofungal richness  
980 and sporocarp production in montado ecosystems. Agrofor. Syst. 82(2), 149-159.

981 Savoie, J.-M.M., Largeteau, M.L., 2011. Production of edible mushrooms in forests: trends in  
982 development of a mycosilviculture. Appl. Microbiol. Biot. 89, 971– 979.

983 Senn-Irlet, B., Bieri, G., 1999. Sporocarp succession of soil-inhabiting macrofungi in an  
984 autochthonous subalpine Norway spruce forest of Switzerland. For. Ecol. Manage. 124(2), 169-175.

985 Shaw, P. J. A., Kibby, G., Mayes, J., 2003. Effects of thinning treatment on an ectomycorrhizal  
986 succession under Scots pine. Mycol. Res. 107(03), 317-328.

987 Sisak, L., Riedl, M., Dudik, R., 2016. Non-market non-timber forest products in the Czech  
988 Republic—Their socio-economic effects and trends in forest land use. Land Use Policy. 50, 390-398.

989 Smith, J.E., Molina, R., Huso, M.M.P., Luoma, D.L., McKay, D., Castellano, M.A., Lebel, T.,  
990 Valachovic, Y., 2002. Species richness, abundance, and composition of hypogeous and epigeous  
991 ectomycorrhizal fungal sporocarps in young, rotation-age, and old-growth stands of Douglas-fir  
992 (*Pseudotsuga menziesii*) in the Cascade Range of Oregon, USA. Can. J. Bot. 80, 186–204.

993 Suz, L.M., Barsoum, N., Benham, S., Dietrich, H.-P., Fetzner, K.D., Fischer, R., García, P., Gehrman,  
994 J., Kristöfel, F., Manninger, M., Neagu, S., Nicolas, M., Oldenburger, J., Raspe, S., Sánchez, G.,  
995 Schröck, H.W., Schubert, A., Verheyen, K., Verstraeten, A., Bidartondo, M.I., 2014. Environmental  
996 drivers of ectomycorrhizal communities in Europe's temperate oak forests. Mol. Ecol. 23 (22), 5628-  
997 5644.

998 Tahvanainen, V., Miina, J., Kurttila, M., Salo, K., 2016. Modelling the yields of marketed mushrooms  
999 in *Picea abies* stands in eastern Finland. For. Ecol. Manage. 362, 79-88.

1000 Taye, Z. M., Martínez-Peña, F., Bonet, J. A., Martínez de Aragón, J., de-Miguel, S., 2016.  
 1001 Meteorological conditions and site characteristics driving edible mushroom production in *Pinus*  
 1002 *pinaster* forests of Central Spain. Fungal Ecol. 23, 30-41.

1003 Tedersoo, L., Suvi, T., Jairus, T., Ostonen, I., Põlme, S., 2009. Revisiting ectomycorrhizal fungi of  
 1004 the genus *Alnus*: differential host specificity, diversity and determinants of the fungal community.  
 1005 New Phytol. 182(3), 727-735.

1006 Tedersoo, L., Bahram, M., Jairus, T., Bechem, E., Chinoya, S., Mpumba, R., Leal, M.,  
 1007 Randrianjohany, E., Razafimandimbison, S., Sadam, A., Naadel, T., Kõljalg, U., 2011. Spatial  
 1008 structure and the effects of host and soil environments on communities of ectomycorrhizal fungi in  
 1009 wooded savannas and rain forests of Continental Africa and Madagascar. Mol. Ecol. 20, 3071–3080.

1010 Urbanová, M., Šnajdr, J., Baldrian, P., 2015. Composition of fungal and bacterial communities in  
 1011 forest litter and soil is largely determined by dominant trees. Soil Biol. Biochem. 84, 53-64.

1012 Vogt, K.A., Bloomfield, J., Ammirati, J.F., Ammirati, S.R., 1992. Sporocarp production by  
 1013 basidiomycetes with emphasis on forest ecosystems. In: Carol, G.C., Wicklow, D.T. (Eds.), The  
 1014 Fungal Community. M. Decker, New York, pp. 563–579.

1015 Wang, Y., Hall, I. R., Evans, L. A., 1997. Ectomycorrhizal fungi with edible fruiting bodies. 1.  
 1016 *Tricholoma matsutake* and related fungi. Economic botany (USA).

1017 Wang, Y., Hall, I.R., 2004. Edible mycorrhizal mushrooms: challenges and achievements. Can. J.  
 1018 Bot. 82:1063–1073.

1019 Wang, Y., Cummings, N., Guerin-Laguette, A., 2012. Cultivation of Basidiomycete edible  
 1020 ectomycorrhizal mushrooms: *Tricholoma*, *Lactarius*, and *Rhizopogon*. In Edible Ectomycorrhizal  
 1021 Mushrooms (pp. 281-304). Springer Berlin Heidelberg.

- 1022 Wang, Y., Chen, Y. L., 2014. Recent Advances in Cultivation of Edible Mycorrhizal Mushrooms. In  
1023 Mycorrhizal Fungi: Use in Sustainable Agriculture and Land Restoration. Pp.: 375-397. Springer  
1024 Berlin Heidelberg.
- 1025 Weigand, J. F., 1997. Forest management for the North American pine mushroom (*Tricholoma*  
1026 *magnivelare* (Peck) Redhead) in the southern Cascade Range (Doctoral dissertation).
- 1027 Wiklund, K., Nilsson, L. O., Jacobsson, S., 1995. Effect of irrigation, fertilization, and artificial  
1028 drought on basidioma production in a Norway spruce stand. Can. J. Bot. 73(2), 200-208.
- 1029 Winder, R. S., Keefer, M. E., 2008. Ecology of the 2004 morel harvest in the Rocky Mountain Forest  
1030 District of British Columbia. Botany 86(10), 1152-1167.
- 1031 Wurtz, T.L., Wiita, A.L., Weber, N.S., Pilz, D., 2005. Harvesting morels after wildfire in Alaska.  
1032 USDA Forest Service Research Note PNW-RN-546.
- 1033 Yamada, A., Maeda, K., Kobayashi, H., Murata, H., 2006. Ectomycorrhizal symbiosis in vitro  
1034 between *Tricholoma matsutake* and *Pinus densiflora* seedlings that resembles naturally occurring  
1035 ‘shiro’. Mycorrhiza. 16(2), 111-116.
- 1036 Zhou, D., Hyde, K. D. 2002. Fungal succession on bamboo in Hong Kong. Fungal Divers. 10: 213-  
1037 227.
- 1038 Zuccherelli, G., 1988. Prime esperienze sulla produzione di piante forestali micorrizate con *Boletus*  
1039 *edulis*. Monti e Boschi. 39, 11–14. (in Italian).
- 1040

1041 Figure 1. Site-specific variables influencing mushroom yield. The inner circle denotes drivers  
1042 typically modified through silviculture and forest management.

1043

1044 Figure 2. Effect of silvicultural operations and forest management practices on mushroom yield.

1045

## Site-specific variables


**Climatic variables**  
(precipitation, temperature, etc.)

**Soil characteristics**  
(pH, soil moisture, etc.)


**Topographical characteristics**  
(slope, aspect, altitude, etc.)

## Stand structure variables

Stand density



Stand age



Tree species composition









POSITIVE EFFECT


NEGATIVE EFFECT



NEGATIVE EFFECT



UNCLEAR (POTENTIALLY POSITIVE)



\*EXCEPT SOME SPECIFIC FIRE-PRONE OR DISTURBANCE-PRONE TAXA

Table 1 Wild epigeous edible mushroom species considered in this review as relevant for mycosilviculture focused on the supply of food provisioning ecosystem services further linked to provision of cultural ecosystems services (e.g., recreation and tourism) and commercial relevance.

Life form	Family	Species	Associated tree/shrub species*	References
ectomycorrhizal	Russulaceae	<i>Lactarius deliciosus</i> s.l. ( <i>Lactarius deliciosus</i> ; <i>Lactarius sanguifluus</i> ; <i>Lactarius vinosus</i> ; <i>Lactarius semisanguifluus</i> )	<i>Pinus sylvestris</i> , <i>Pinus nigra</i> , <i>Pinus halepensis</i> , <i>Pinus pinaster</i> , <i>Pinus contorta</i> , <i>Pinus radiata</i> , <i>Pseudotsuga menziesii</i>	Guinberteau et al., 1990; Smith et al., 2002; Martínez de Aragón et al., 2007; Ortega-Martínez et al., 2011; Bonet et al., 2012; Martínez-Peña et al., 2012a; Martínez-Peña et al., 2012b; Ágreda et al., 2014; de-Miguel et al., 2014; Guerin-Laguette et al. 2014; Kucuker and Baskent, 2015; Liu et al., 2016
		<i>Lactarius deterrimus</i>	<i>Picea</i> spp.	Buée et al., 2011
		<i>Lactarius salmonicolor</i>	<i>Abies alba</i> , <i>Abies bormülleriana</i>	Courtecuisse and Duhem, 1994; Kucuker and Baskent, 2015
		<i>Lactarius volemus</i>	<i>Pinus</i> spp.	Liu et al., 2007
	Boletaceae	<i>Boletus edulis</i> s.l. ( <i>Boletus aestivalis</i> , <i>Boletus aereus</i> , <i>Boletus edulis</i> , <i>Boletus pinophilus</i> )	<i>Abies alba</i> , <i>Cistus ladanifer</i> , <i>Quercus</i> spp., <i>Picea</i> spp., <i>Pinus sylvestris</i>	Nocentini et al., 2004; Salerni and Perini, 2004; Oriade-Rueda et al., 2008; Ortega-Martínez et al., 2011; Martínez-Peña et al., 2012b; de-Miguel et al., 2014; Hernández-Rodríguez et al., 2015a, Hernández-Rodríguez et al., 2015b; Mediavilla et al., 2016; Tahvanainen et al., 2016; Parladé et al. 2017
		<i>Suillus granulatus</i>	<i>Pinus</i> spp.	Savoie and Largetau, 2011

		<i>Suillus bovinus</i>	<i>Pinus sylvestris</i>	Shaw et al., 2003
		<i>Suillus brevipes, Suillus tomentosus</i>	<i>Pinus contorta</i>	Kropp and Albee, 1996
		<i>Leccinum corsicum</i>	<i>Cistus ladanifer</i>	Martín-Pinto et al., 2006
	<i>Tricholomataceae</i>	<i>Tricholoma magnivelare,</i>	<i>Abies amabilis, Abies magnifica var. shastensis, Pinus contorta, Pinus monticola, Pseudotsuga menziesii, Tsuga heterophylla, Tsuga mertensiana</i>	Weigand, 1997, Kranabetter et al., 2002; Kranabetter et al., 2005; Luoma et al., 2006;
		<i>Tricholoma matsutake</i>	<i>Pinus densiflora</i>	Wang et al., 1997; Wang and Hall, 2004; Wang et al., 2012; Wang and Chen, 2014; Yamada et al., 2006
	<i>Cantharellaceae</i>	<i>Cantharellus cibarius</i>	<i>Picea excelsa</i>	Danell et al., 2002, Wang and Chen, 2014
		<i>Cantharellus formosus</i>	<i>Picea sitchensis, Pseudotsuga menziesii, Tsuga heterophylla</i>	Kranabetter et al. 2009; Pilz et al. 2006
saprotrophic and ectomycorrhizal**	<i>Morchellaceae</i>	<i>Morchella spp. (Morchella conica s.l.; Morchella esculenta s.l.)</i>	<i>Abies concolor, Abies grandis, Abies lasiocarpa, Asimina triloba, Larix occidentalis, Picea glauca, Pinus contorta, Pinus lambertiana, Pinus ponderosa, Pseudotsuga menziesii, Tilia Americana, Ulmus americana</i>	Pilz et al., 2004; Mihail et al., 2007; Pilz et al., 2007; Winder and Keefer, 2008; Larson et al., 2016
saprotrophic and parasitic	<i>Physalacriaceae</i>	<i>Armillaria mellea s.l. (Armillaria mellea, Armillaria borealis, Armillaria cepistipes, Armillaria gallica,</i>	<i>Abies grandis, Clintonia uniflora, Physocarpus malvaceus, Quercus spp., Thuja plicata, Tsuga heterophylla</i>	Kim et al., 2010; Lee et al., 2016

		<i>Armillaria ostoyae</i> , <i>Armillaria tabescens</i> )		
--	--	--	--	--

\* the species listed here are those characterizing study areas of cited mycosilviculture researches

\*\* Morchella spp. can behave either as saprotrophic or mycorrhizal at different stages of their life cycle (Pilz et al., 2007).

Table 2. State of knowledge about major research topics examined in the present review concerning relationship between forest stand attributes, forest management practices, main disturbances and mushroom occurrence. Symbols indicate current knowledge about the topic: “absence or very little knowledge present in the literature” (--); “little knowledge and few experiences present in the literature” (-); “several studies dealing with the topic, but further research is still needed” (+).

Forest stand age	Influence of forest stand age on wild mushroom production	+
	Mushroom production in even-aged forests	+
	Mushroom production in uneven-aged forests	-
Stand density	Relationship between basal area and wild mushroom production	+
	Relationship between n° of trees per hectare and wild mushroom production	-
	Relationship between canopy cover and wild mushroom production	+
Tree species composition	Influence of tree species composition and diversity on wild mushroom production	--
	Mushroom production in coniferous forests	+
	Mushroom production in broadleaved forests	-
Effect of forest management practices	Effect of thinning on wild mushrooms production in the short term	+
	Effect of thinning on wild mushrooms production in the long term	-
	Effect of clear-cuts on wild mushrooms production	+
	Effect of shelterwood methods on wild mushrooms production	-
	Effect of selective cutting on wild mushrooms production	-
	Use of mycorrhized trees for enhancing mushroom production in natural forest ecosystems	--
	Effect of timber harvesting on wild mushrooms production	-

	Effect of shrub control on wild mushrooms production	--
	Absence of forest management	--
Effect of other disturbances	Effect of picking activity on wild mushrooms production	+
	Effect of removal of litter on wild mushrooms production	-
	Effect of fire on wild mushrooms production	+